Armed Services Technical Information Agency

Reproduced DOCUMENT SERVICE CENTER KNOTT BUILDING, DAYTON, 2, OHIO

This document is the property of the United States Government. It is furnished for the duration of the contract and shall be returned when no longer required, or upon recall by ASTIA to the following address: Armed Services Technical Information Agency, Document Service Center, Knott Building, Dayton 2, Ohio.

NOTICE: WHEN GOVERNMENT OR OTHER DRAWINGS, SPECIFICATIONS OR OTHER DATA ARE USED FOR ANY PURPOSE OTHER THAN IN CONNECTION WITH A DEFINITELY RELATED GOVERNMENT PROCUREMENT OPERATION, THE U. S. GOVERNMENT THEREBY INCURS NO RESPONSIBILITY, NOR ANY OBLIGATION WHATSOEVER; AND THE FACT THAT THE GOVERNMENT MAY HAVE FORMULATED, FURNISHED, OR IN ANY WAY SUPPLIED THE SAID DRAWINGS, SPECIFICATIONS, OR OTHER DATA IS NOT TO BE REGARDED BY IMPLICATION OR OTHERWISE AS IN ANY MANNER LICENSING THE HOLDER OR ANY OTHER PERSON OR CORPORATION, OR CONVEYING ANY RIGHTS OR PERMISSION TO MANUFACTURE. USE OR SELL ANY PATENTED INVENTION THAT MAY IN ANY WAY BE RELATED THERETO.

BEST

AVAILABLE

COPY



FCPAC

Studies of the Rare-Earth lydrides Technical Report X

DISCUSSION AND SUMMARI

Office of Naval Research Physical Sciences Division

Project No. NR 356-290

Contract No. Nonr 228(03)

Ву

James C. Warf, Project Supervisor William L. Korst, Research Assistant

The Department of Chemistry University of Southern California Los Angeles 7, California

June 30, 1956

Title List of Technical Reports on "Studies of the Bre-Earth Hydrides" June 30, 1956

Office of Naval Research Physical Sciences Division

Project No. NR 356-290

Contract No. Nonr 228(03)

рÀ

James C. Warf, Project Supervisor and Milliam L. Korst, Research Assistant The Department of Chemistry University of Southern California

Los Angeles 7, Galifornia

Technical Report I. A Survey of the Literature concerning the Rare-Earth Hydrides.

Technical Report II. Dispersions of Rame-Earth Hydrides in Mercury. High-Temperature Thermostatted Furnace. Silica Torsion Balance.

Technical ReportIII. Pressure-Temperature-Composition Studies of the Lanthanum-,
Cerium-, Prassodymium-, Neodymium-, Samarium-, and
Ttterbium-Hydrogen Systems. Experimental.

Technical Report IV. Pressure-Temperature-Composition Studies of the Lanthanum-, Cerium-, Praseodymium-, Neodymium-, Samarium-, and Itterbium-Hydrogen Systems. Results.

Technical Report V. The Crystal Structure of the Deuterides of Ytterbium and Europium.

Technical Report VI. Solutions of Europium and Ytterbium Metals in Liquid Ammonia

Technical Report VII. Crystal Structure of the Rare-Earth Metals.

Technical Report WII. Crystal Structure of the Rare-Earth Hydrides. X-Ray Diffraction by Rare-Earth Hydride Amalgums.

Technical Report IX. Lanthanum Monoxide.

Technical Report X. Discussion and Summary.

DISCUSSION AND SUMMARY

Discussion

The results obtained in the different fields of investigation of the various rare earth metal-hydrogen systems have already been discussed in the individual chapters of this dissertation. Evidence has been presented supporting a plausible explanation of the nature and constitution of the rare earth hydrides. This explanation may at best be called a hypothesis; additional experimental studies, such as magnetic measurements, and measurements of electrical conductivity of hydrides of various compositions, are needed for further elucidation of their constitution.

Summary

A physical chemical investigation of the hydrides of various rare earth metals has been carried out. The principal fields of investigation have been studies of pressure-temperature-composition relationships, and X-ray diffraction studies of the powdered hydrides.

The properties and behavior of the hydrides formed by lanthanum, cerium, praseodymium, and neodymium are very similar. Dissociation pressure isotherms for these systems

consist of three branches: 1) an initial ascending branch at low hydrogen composition, indicative of the solution of hydrogen in the metal to form a single solid phase; 2) a constant-pressure plateau indicating the co-existence of two solid phases; 3) a final, steeply ascending curve extending into the region of higher hydrogen composition. The relative dissociation pressures of the hydrides and deuterides of lanthanum and cerium have been compared and discussed.

x-ray diffraction studies have confirmed the facecentered cubic structures of these hydrides, which are
greatly expanded over the structures of the metals. This
expansion is a maximum for the hydrogen-rich phase of the
two-phase region; with further addition of hydrogen the
lattice contracts slightly.

Similarities between the hydrides and the monochalcogenides of these rare earth elements have been pointed
out, and the probable nature of the bonding in the hydrides
has been discussed.

The properties of dispersions of the hydrides of lanthanum and cerium in mercury have been investigated.

The metals europium and ytterbium have been found to form dihydrides which are isostructural with the hydrides of the alkaline earth metals. These two rare earth metals

have also been found to dissolve in liquid ammonia. Some of the properties of these metal-in-ammonia solutions have been investigated.

The structures of the rare earth metals themselves have also been investigated, and evidence has been found showing that the hexagonal modifications of the metals do not possess a simple hexagonal closs-packed structure.

Evidence for the existence of lanthanum monoxide has been obtained; indications of similar compounds of other rare earth metals have also been found.

See Technical Report 1 for references